

Full Length Research Paper

# Comparison the chemical components and functional properties of fish protein isolate (FPI) from *Pangasius hypophthalmus* byproducts to ones of other Protein Isolates (PIs)

\*<sup>1</sup>Thuy C.X, <sup>2</sup>Lam T.B, <sup>3</sup>Toan H.T, <sup>4</sup>Commick K.Mc

<sup>1</sup>Faculty of Food Technology, Hochiminh City University of Food Industry (HUFI)- Vietnam

<sup>2</sup>Department of Food Technology, HCMC University of Technology (VNU) - Vietnam

<sup>3</sup>Biotechnology R and D Institute, Can Tho University (CTU) - Vietnam

<sup>4</sup>Minnesota University, USA

Corresponding Author's E-mail: thuycx@cntp.edu.vn

Accepted 22<sup>nd</sup> April, 2014

The researching on chemical composition and foaming, emulsifying abilities of fish protein isolate (FPI) which obtained in the hydrolysis of *Pangasius hypophthalmus* byproducts were proceeded; and compared to the ones of commercial soy protein isolate (SPI), commercial whey protein isolate (WPI). The results showed that the FPI's foaming ability was equivalent to WPI's one and higher than SPI's one. The highest foaming abilities of FPI, WPI, SPI were  $94.61 \pm 1.03\%$  (at pH=7);  $96.42 \pm 1.12\%$  (at pH=7);  $80.54 \pm 0.89\%$  (at pH=8) respectively. Emulsifying abilities of FPI and SPI were equal and both reached the highest values at pH=7. The maximum emulsifying ability of FPI was  $21.03 \pm 1.01$  mL oil/g FPI while the highest one of SPI was  $21.56 \pm 0.91$  mL oil/g SPI. Emulsifying ability of WPI was lower than the ones of FPI, SPI. The protein component in FPI, SPI and WPI was higher than 90%. The very low lipid contents in FPI, WPI, SPI were  $0.94 \pm 0.18\%$ ;  $0.81 \pm 0.05\%$ ;  $0.39 \pm 0.08\%$  respectively. Moisture and ash contents of the FPI, WPI, SPI were  $2.86 \pm 0.90\%$  and  $4.94 \pm 0.16\%$ ;  $3.01 \pm 0.02\%$  and  $5.17 \pm 0.06\%$ ;  $4.18 \pm 0.42$  and  $4.45 \pm 0.24\%$  respectively.

**Keywords:** Fish protein isolate, *Pangasius hypophthalmus* byproducts, FPI functional properties, PIs functional properties

## INTRODUCTION

Improving the functional properties of protein products (protein hydrolysates - PHs; protein concentrates - PCs; protein isolates - PIs), including: solubility, water holding, oil holding, emulsifying, and foaming characteristics are a major challenge for food science (I. R. Freitas, 2011). The protein products may be used as food ingredients or food additives due to their functional properties (Liceaga-Gesualdo and Li-Chan, 1999). The use of fish waste has been increasing interests in years. It is considered to be a safe, high-protein material with many nutritional benefits (Guerard et al., 2002). FPI contains proteins with small molecular weight and can be called peptone from fish (Asbjorn Gildberg, 2007). FPI's functional properties as well as biological activities depend on its origin and produced methods (Samanta S. Khora, 2013). For Vietnam's *Pangasius hypophthalmus* byproducts,

under controlled conditions, enzymatic hydrolysis influences the molecular weight, hydrophobicity, and polar groups of the proteins in final products (Hoa et al., 2012). The characteristics of the proteins in FPI directly affect its functional properties, such as emulsifying and foaming abilities (Kristinsson et al., 2000; Gbogouri et al., 2004). Both of whey and soy proteins are by-products of the industry. The functional properties of WPI and SPI, mainly among them were solubility, emulsifying, gelling and foaming abilities (H.E. Swaiswood, 1996; L.M. Huffman, 1998). Commercial SPI was manufactured from defatted soy flakes by separation of the soy proteins from both the soluble and the insoluble carbohydrate fractions of the soybean. WPI is obtained by removing sufficient non-protein constituents from whey so that the finished dry product

## 2. Glob. J. Food Sci. Technol.

contains no less than 90% protein (Burrington, 2000). The whey's fat was first removed by micro-filtration (MF) and then ultra-filtration (UF) or nano-filtration (NF). In addition to concentrating protein and fractionating whey into individual proteins, WPI can be subjected to controlled enzyme hydrolysis in order to yield smaller protein fragments (Russell Tara Alexandra, 2004).

There are many studies on the foaming, emulsifying properties of WPI (El-Shiniby et al., 2007; I. Nicorescu et al., 2011; Panizzolo et al., 2012...); SPI (Rickert, 2004; Deak, 2007; Egbert, 2004...) and some studies about FPI's ones (Rong et al., 2011; San et al., 2008...). But there no comparative studies among them. Therefore, the objective of this study is to conduct a comparative examination for the foaming and emulsifying abilities as well as chemical components of FPI to the ones of commercial WPI, SPI.

## MATERIALS AND METHODS

### Material

*By-products (the spine and head...)* of *Pangasius hypophthalmus* were received from Can Tho Fish Joint Stock Company (CAFICO) - Mekong River Delta, Vietnam. Then it had been refrigerated, transported to the laboratory, divided into small unit for each experiment and stored at  $-20^{\circ}\text{C}$  until used.

*Enzymes Alcalase 2.4L* was purchased from EAC Co., Ltd. (sole-exclusive agent for Novozyme in Ho Chi Minh city, Vietnam).

*Commercial SPI (SPI<sub>4</sub>)* was purchased from Prestige L.O. Limited (France). According to brochure Prestige L.O. Limited: SPI was obtained by removing soluble carbohydrates, defatted soy meal by using aqueous or alkali extraction of proteins at a pH range of 7-10; dispersion of the precipitate on alkaline medium (pH 8.0), further processing by ultra-filtration and freeze-dried to get SPI powder.

*Commercial WPI* was purchased from Labrada Nutrition (Toronto, Canada). According to brochure of Labrada Nutrition: WPI was obtained by removing sufficient non-protein constituents from whey; fat was first removed by cooling, then microfiltration and ultra-filtration or nano-filtration; free-dried to get WPI powder.

All chemical reagents used for the experiments were in analytical grade.

### Methods

#### Hydrolysis process and collection FPI

Hydrolyzing the *Pangasius hypophthalmus* by-products by protease (Alcalase 2.4L) under controlled condition.

After hydrolysis, filtering to separate the solid and liquid, inactivating enzyme Alcalase 2.4L by heat treatment at  $90^{\circ}\text{C}/10$  minutes as recommendation of Novozymes. Hydrolyzed solution was then cooled to  $4^{\circ}\text{C}$  for a preliminary de-fatting, vacuum filtered through non-ash paper and then centrifuged to de-fat at the speed of 15,000 rpm for 20 minutes.

The solution obtained after centrifugation was brought to freeze-dry to get FPI powder. FPI powder is used to study the foaming and emulsifying abilities.

#### Chemical analysis of PIs

The moisture and ash content were determined according to the AOAC standard methods 930.15 and 942.05 respectively. Total nitrogen content of FPIs was determined by using the Kjeldahl method. Lipids were determined gravimetrically after Soxhlet extraction of dried samples with hexane. All measurements were performed in triplicate.

#### Determination of PI's foaming ability

Hydrolyzing *Pangasius hypophthalmus* by-products by protease (Alcalase 2.4L) under controlled conditions to get FPI with highest foaming ability as follows: enzyme/substrate (E/S) ratio of 0.2% (v/w); hydrolysis temperature is  $64^{\circ}\text{C}$ ; hydrolysis time is 92 minutes. After collecting the FPI powder, comparing foaming ability of FPI to the ones of WPI, SPI.

Foaming ability of PIs was determined by the method of Kazunobu et al (2005) and Watanabe et al. (1981): 0.25 g FPI would be dissolved in 25 ml of distilled water. The mixture was adjusted to pH 4.0, 5.0, 6.0, 7.0, 8.0 and 9.0 by 0.5N NaOH or HCl. Then it was stirred by electric mixer to create foam system at room temperature. The sample after stirring was poured into the instrument (flash) for measuring both the total volume in foaming phase and the volume of separated water after 30 seconds. Foaming ability is calculated as follows:

$$\text{FA}(\%) = \frac{V_f - V_w}{V_i} * 100$$

Where:  $V_f$ : total volume in foaming phase;  $V_w$ : volume of separated water;  $V_i$ : volume of initial mixture

#### Determination of PI's emulsifying ability

Hydrolyzing *Pangasius hypophthalmus* by-products by protease (Alcalase 2.4L) under controlled conditions to get FPI with highest emulsifying ability as follows: pH 7.4; E/S ratio is 0.19% (w/v), temperature at  $62^{\circ}\text{C}$ , hydrolysis time is 80 minutes. After collecting the FPI

**Table 1.** Chemical composition<sup>(\*)</sup> of the PIs

Source	Protein (%)	Lipid (%)	Moisture (%)	Ash (%)
FPI	90.14 ± 1.60 <sup>a</sup>	0.94 ± 0.18 <sup>a</sup>	2.86 ± 0.90 <sup>a</sup>	4.94 ± 0.16 <sup>a</sup>
Commercial WPI	90.74 ± 0.84 <sup>a</sup>	0.81 ± 0.05 <sup>a</sup>	3.01 ± 0.02 <sup>b</sup>	5.17 ± 0.06 <sup>b</sup>
Commercial SPI	90.08 ± 1.43 <sup>a</sup>	0.39 ± 0.08 <sup>b</sup>	4.18 ± 0.42 <sup>c</sup>	4.45 ± 0.24 <sup>c</sup>

(\*) Results reported are means of triplicate samples ± standard deviation. Values in the same column with different superscripts are significant different at p<0.05

powder, comparing emulsifying ability of FPI to the ones of WPI, SPI

Emulsifying capacity of PIs was measured as described by Rakesh and Metz (1973), with some modification. One gram of each freeze-dried sample was transferred into a 250 mL beaker and dissolved in 50 mL of 0.5 N NaCl and then 50 mL of soybeans pure oil was added. The solution was adjusted to pH of 4.0, 5.0, 6.0, 7.0, 8.0 and 9.0 by 0.5N NaOH or HCl. Homogenizing the solution for 120 sec. at 10.000 rpm to make an emulsion. The mixture was transferred into centrifuge tubes, kept under a water-bath at 90°C for 10 min and then centrifuged at 3000 rpm for 20 min. Emulsifying capacity was calculated using the equation:

$$EC \text{ (mL oil/g FPI)} = (V_A - V_R) / W_S$$

Where:  $V_A$  is the volume of oil added to form an emulsion;  $V_R$  is the volume of oil released after centrifugation;  $W_S$  is the weight of the sample.

### Statistical analysis

All analytical determinations were carried out in triplicate and mean values with standard deviation (SD) are presented. Results were analyzed statistically by ANOVA using SPSS 15.0 to ascertain whether differences were significant at p<0.05.

## RESULTS AND DISCUSSION

### Chemical composition of PIs

The chemical composition of FPI from *Pangasius hypophthalmus*, commercial WPI, commercial SPI was determined. The results were shown in Table 1.

Base on above results in Table 1, the protein content of all three PIs was higher than 90% and have no significant difference (p <0.05). Protein content in the aforementioned PIs is similar. High protein content reflected the quality of the PIs. Moisture and ash contents in WPI, FPI, SPI were different (p <0.05). The FPI's moisture is lowest while SPI's one is highest. Ash content of WPI is highest and the lowest one belongs to SPI.

About FPI from *Pangasius hypophthalmus*, our studying results had been similar to the findings of other investigators whom reported protein content ranging

from 78% to 93% for lyophilized hydrolysate or FPI samples from *Pollachius virens* (Gholam et al., 2012); *Catla catla* (Balaswamy et al., 2011); *Salmon* (Kristinson et al., 2000), and *Pacific whiting* muscle (Pacheco-Aguilar, 2008). Ash and moisture contents in FPI from *Pangasius hypophthalmus* byproducts were equal to ones of the FPI from *Silver catfish* (3.99% ÷ 5.61% and 3.33% ÷ 4.45% respectively) (Azima et al., 2013).

The fat content in all 3 types of PIs was very low (all less than 1%). The SPI's fat content was lowest in comparison with WPI, FPI. The fat content in WPI and FPI was similar and have no significant difference (p <0.05). Lipid content in FPI and WPI was higher than the one in SPI because both WPI and FPI derived from animals while SPI derived from vegetable (Russell Tara Alexandra, 2004). The lipid in FPI was highest due to *Pangasius hypophthalmus* belonged to fat catfish group. The lipid content in *Pangasius hypophthalmus* byproducts was 32.21±1.89% (Hoa, 2012). In generally, PIs which obtained from different production methods, the chemical compositions were almost similar but functional properties could be very different (these will be examined in the following section).

### Emulsifying ability of PIs

Based on the results presented in Figure 1, WPI's emulsifying ability is lowest and there is significant difference (p <0.05) compared with emulsifying ability of SPI, FPI. Emulsifying abilities of FPI and SPI are similar and have no significant differences (p <0.05). Emulsifying ability of all WPI, SPI, FPI have reached the lowest value at lightly acid pH (pH=5). The highest emulsifying capabilities of SPI, FPI and WPI were obtained at pH=7.0. The maximum emulsifying ability of WPI compared with the maximum ones of FPI, SPI was only 83.95% and 86.07% respectively. At pH values which were lower than 7.0, the emulsifying ability of PIs was low. In contrast, at the pH values of 7.0 or higher, emulsifying ability of PIs achieved maximum values, and then declined lightly.

This is explained as follows: a significant increasing in emulsifying capacity of PIs at pH=7.0 may be due to higher quantities of soluble proteins in PIs (Ann Elizabeth Theodore. 2005). The pH also affects emulsifying property by changing the solubility and surface hydrophobicity of proteins, as well as the charge




Figure 1. The emulsifying ability of WPI, FPI, SPI




Figure 2. The foaming ability of WPI, FPI, SPI

of the protective layer surrounding the lipid globules. Ions alter the electrostatic interactions, conformation, solubility of the proteins, and hydrophilic-lipophilic balance (Sikorski, 2002). WPI's emulsifying ability was lower than the ones of FPI, SPI because as the number of medium and small size proteins (7-20 kDa) in FPI and SPI are similar and accounted for 65 ÷ 75% of their protein content while this WPI's size of protein group was only 35.8%. This group of proteins has an important role in forming the emulsifying ability of PIs (Cecilia Abirached et al., 2012; IR Freitas et al., 2011; Ameri Shahrabi A et al., 2011). On the other hand, environmental pH also affects emulsifying properties by changing the solubility and surface hydrophobicity of proteins, as well as the charge of the protective layer surrounding the lipid globules. Ions alter the electrostatic interactions, conformation, solubility of the proteins, and hydrophilic-lipophilic balance (Sikorski, 2002). At high pH values, the emulsifying ability of PIs increased because at these pH values, small and medium polypeptides (from 7-20 kDa) can be unfolded due to negative charges. Repulsion could be resulted from this change and allowing for better orientation at the interface (Pacheco-Aguilar et al., 2008). This could result in a

more efficient exposure of hydrophilic and hydrophobic residues in these peptides, promoting a major interaction at the oil-water (O:W) interface. Since the lowest solubility occurred at pH 4.0, 5.0 and 6.0, peptides could not move rapidly to the interface. Additionally, the net charge of the peptide will be minimized at these pH values. So the emulsifying ability of PIs decreased.

Our results is similar to what reported from Taheri A., 2011 about emulsifying ability of FPI from rainbow trout (*Onchorhynchus mykiss*) viscera; Klompong et al., 2007 about emulsifying ability of FPI from *yellow-striped trevally*; Mohamed Beva Kelfala Foh, 2012 about emulsifying ability of FPI from Tilapia (*Oreochromis niloticus*); Russell Tara Alexandra, 2004 about emulsifying ability of WPI, SPI.

#### Foaming capability of PIs

The foaming properties of the 3 PIs were determined at pH values of 4.0, 5.0, 6.0, 7.0, 8.0 and 9.0. The foaming capacity was shown in Figure 2.

The foaming capacity of FPI, WPI, and SPI ranged between  $58.21 \pm 0.78\%$  ÷  $94.61 \pm 1.03\%$ ;  $55.46 \pm 0.09$  ÷

96.42±1.12%; and 38.26±0.47%÷ 80.54±0.98% respectively. The foaming capacities of all 3 types of PIs have reached the lowest value at pH of 4.0 ÷ 5.0; when pH increases, the foaming ability of WPI, SPI, FPI tends to increase up to a maximum at pH of 7.0 ÷ 8.0. At neutral to alkaline pH, the foaming ability of FPI, SPI, and WPI was higher than the one at light acid pH. Foaming ability of WPI and FPI was similar and much higher than SPI's. The highest foaming ability of WPI, FPI reached at pH=7.0 and have no statistically significance ( $p < 0.05$ ). The highest foaming ability of SPI was only 83.50% and 85.13% compared with WPI and FPI respectively.

This is explained as follows: Foaming ability is related to decreasing's rate of the surface tension of the air/water interface caused by absorption of protein molecules (Sathe et al., 1982). Good foaming ability was linked with flexible protein molecules, which reduces surface tension. Low foaming ability on the other hand can be related to highly ordered globular proteins, which resists surface denaturation. The basic requirements of proteins as good foaming agents are the ability to: (1) absorb the proteins rapidly at air water interface during bubbling, (2) undergo rapidly conformational change and rearrangement at the interface, and (3) form a cohesive viscoelastic film via intermolecular interactions. The first two factors are essential for better foaming ability whereas the third is important for the stability of the foam (Sathe et al., 1982; Graham and Phillips, 1976; Russell, 2004; Anusha, 2010). Although the total protein in WPI, FPI, SPI is similar; the ratio of proteins in the same molecular weight groups is different. This will affect the foaming ability of each PIs (Mohamed et al., 2012; Ann, 2005). It should be noted that the adsorption rate to the air-water interface may be influenced by the molecular size, protein structure and hydrophobicity of the hydrolysates (Martin et al., 2002). These are highly dependent on both producing methods of PIs and the parent protein from which they are obtained and the hydrolysis procedure. The hydrolysis of protein produces a range of peptides that possess altered hydrophobicity, net charge, and conformation in comparison to the native molecule. Their reduced molecular weight makes them more flexible, form a stable interfacial layer and increase the rate of diffusion to the interface, which in turn improves foaming ability (Wilde and Clark, 1996). The foaming ability of all WPI, FPI, SPI are low at low pH due to the lowest foaming ability was attributed to the protein behavior at around its isoelectric point. At high pH, it was likely due to the increased net charges on the protein, which weakened the hydrophobic interactions but increased the flexibility of the protein. This allowed the protein to diffuse more rapidly to the air-water interface to encapsulate air particles and then enhance the foam formation (Wierenga and Gruppen 2010).

Our study results were equivalent to the previous ones that have been reported. SPI's foaming ability is less than foaming ability of WPI and SPI (Cecilia et al., 2012; Freitas et al., 2011). Foaming ability of FPI from Tilapia

(*Oreochromis niloticus*) could reach 89.63% while SPI's one is only 71.52% at pH 7.0 ÷ 8.0 (Mohamed et al., 2012). The highest foaming ability of SPI from defatted soy is about 80% (Ameri et al., 2011). The highest foaming ability of FPI from Sardinella (*Sardinella aurita*) reached 92.8% at pH=7 (Nabil Souissi et al., 2007).

## CONCLUSION

Protein isolate obtained from enzymatic hydrolysis of *Pangasius hypophthalmus* byproducts has relatively good abilities of foaming and emulsifying. FPI's foaming ability was equivalent to WPI's one and higher than the one of SPI. Emulsifying ability of FPI from *Pangasius hypophthalmus* byproducts was equivalent to SPI's one and higher than the one of WPI. Highest foaming ability of FPI from *Pangasius hypophthalmus* byproducts and WPI were 94.61±1.03% and 96.42±1.12% (at pH=7.0) respectively. The lowest FPI's foaming ability was 58.21±0.78% (at pH=5.0) while the one of WPI was 55.46±0.09 (at pH=4.0). The highest emulsifying ability of FPI was 21.03±1.01mL oil/g FPI (at pH=7) and the one of SPI was 21.56±0.91mL oil/g SPI (at pH 7). The lowest emulsifying ability of FPI and SPI (both at pH=5) are 10.11±0.26 mL oil/g FPI and 11.32±0.62 mL oil/g SPI respectively. The protein content in FPI, SPI, WPI was similar and over 90%. Fat content in FPI, WPI and SPI was 0.94±0.18%; 0.81±0.05% and 0.39±0.08% respectively. This was very low fat amount (less than 1%). Other chemical compositions, such as: moisture and ash content of the FPI, WPI, SPI were 2.86±0.90% and 4.94±0.16%; 3.01±0.02% and 5.17±0.06%; 4.18±0.42 and 4.45±0.24% respectively.

## REFERENCES

- Ameri SA, Badii F, Ehsani MR, Maftoonazad N, Sarmadizadeh D (2011). Functional and thermal properties of chickpea and soy-protein concentrates and isolates. Iranian Food J. 38:167-174.
- Ann ET(2005). Bioactive and functional properties of catfish protein hydrolysates and catfish protein isolates. A thesis for the degree of MSc. of science, Uni. of Florida.
- Anusha GPS (2010). Pacific Hake (*Merluccius productus*) fish protein hydrolysates with antioxidative properties. A thesis for the degree of Doctor of Philosophy in the faculty of graduate studies (Food Science), The University of British Columbia.
- Asbjorn G (2007). Enzymes and Bioactive Peptides from Fish Waste Related to Fish Silage, Fish Feed and Fish Sauce Production. Haworth Press. Inc: 1255-1262.
- Amiza MAYW, Faazaz AL (2013). Physicochemical properties of silver catfish (*Pangasius sp.*) frame hydrolysate. Int. Food Res. J. 20(3):1255-1262.
- Balaswamy K, Prabhakara Rao PG, Narsing Rao G, Jyothirmayi T (2011). Functional properties of roe protein hydrolysates from *Catla Catla*. Electr. J. Environ., Agric. Food Chem. 10(4):2139-2147.

- Burrington K (2000). Nutritional and Beneficial Ingredients. Food Product Design 3000: 1-12.
- Cecilia Abirached, Claudia Alejandra Medrano, Ana Claudia Araujo, Patrick Moyna, and Luis Alberto Panizzolo (2012). Comparison of Interfacial and Foaming Properties of Soy and Whey Protein Isolates. J. Food Sci. Engine. 2:376-381.
- Deak NA, Murphy PA, Johnson LA (2007). Characterization of fractionated soy proteins produced by a new simplified procedure. J. Am. Oil Chem. Soc. 84:137-149.
- El-Shiniby S, Farrag AF, El-Garawany G, Assem FM (2007). Rheological and functional properties of whey protein concentrate and  $\beta$ -lactoglobulin and  $\alpha$ -lactalbumin rich fractions. Int. J. Dairy Sci. 2:196-206.
- Freitas IR, GV, Rios DG, Prentice C (2011). Functionality of Protein Isolates from Argentine Anchovy (*Engraulis anchoita*) Residue Obtained Using pH Shift Processing. J. Food Sci. Engine. 1:374-378
- Gbogouri GA, Linder M, Fanni J, Parmentier M (2004). Influence of hydrolysis degree on the functional properties of salmon byproduct hydrolysates. J. Food Sci. 69:615–622.
- Gholam Reza Shaviklo (2008). Evaluation and Utilisation of Fish Protein Isolate Products. Master of Science in Food Science, University of Iceland.
- Graham DE, Phillips MC (1976). Foams. Academic Press, London, pp 237.
- Guerard F, Guimas L, Binet A (2002). Production of tuna waste hydrolysates by a commercial neutral protease preparation. J. Mole. Catalysis B: Enzymatic 19-20:489-498.
- Huffman LM (1998). The importance of whey protein fractions for WPC and WPI functionality, in: Whey. Proceedings of the Second International Whey Conference, pp. 197-205.
- Hoa MX, Lam TB (2008). Optimization of enzymatic hydrolysis of viscera of Pangasiidae to obtain calcium-binding protein hydrolysate. Vietnam Journal of Agriculture and Rural development 38: 124-132.
- Kazunobu Tsumuraa et al. (2005). Functional properties of soy protein hydrolysates obtained by selective proteolysis. Food science and Technology 38:255 - 261.
- Klompong V, Benjakul S, Kantachote D, Shahidi F (2007). Antioxidative activity and functional properties of protein hydrolysate of yellow stripe trevally (*Selaroides leptolepis*) as influenced by the degree of hydrolysis and enzyme type. Food Chemistry 102:120-131.
- Kristinsson HG, Rasco BA (2000). Hydrolysis of salmon muscle proteins by an enzyme mixture extracted from Atlantic salmon pyloric caeca. Journal of Food Biochemistry 24: 177-187.
- Liceaga-Gesualdo AM, Li-Chan ECY (1999). Functional Properties of Fish Protein Hydrolysate from Herring (*Clupea harengus*). Journal of Food Science 64(6): 1000-1004.
- Martin AH, Grolle K, Bos MA, Stuart MAC and VanVliet T (2002). Network forming properties of various proteins adsorbed at the air/water interface in relation to foam stability. J. Colloid and Interface Sci. 254:175-183.
- Mohamed BKF, Xia W, Issoufou A, Qixing J (2012). Influence of pH Shift on Functional Properties of Protein Isolated of Tilapia (*Oreochromis niloticus*) Muscles and of Soy Protein Isolate. Food Bioprocess Technol 5:2192-2200.
- Nabil S, Ali B, Yousra T-E, Moncef N (2007). Biochemical and Functional Properties of *Sardinella* (*Sardinella aurita*) By-Product Hydrolysates. Food Technol. Biotechnol. 45(2):187-194.
- Nicorescu I, Vial C, Talansier E, Lechevalier V, Loisel C, Della Valle D (2011). Comparative effect of thermal treatment on the physicochemical properties of whey and egg white protein foams, Food Hydrocolloids 25:797-808.
- Pacheco-Aguilar R, Mazorra-Manzano MA, Ramírez-Suárez JC (2008). Functional properties of fish protein hydrolysates from Pacific whiting (*Merluccius productus*) muscle produced by a commercial protease. Food Chemistry 109:782-789
- Panizzolo LA, Mussio LE, Añón MC (2012). A kinetic description for the destabilization process of protein foams. Int. J. Food Properties 15:60-68.
- Rickert DA, Johnson LA, Murphy PA (2004). Functional properties of improved Glycinin and  $\beta$ -conglycinin fractions. J. food Sci. 69:303-311.
- Rakesh J, Metz A (1973). Acid precipitated fish Protein isolate exhibits good functional properties. Food Product Development 7:18-24.
- Rong L, Hari BK, Wentong X, Chuyi L (2011). Characterization of Allergens Isolated from the Freshwater Fish Blunt Snout Bream (*Megalobrama amblycephala*). J. Agric. Food Chem. 59:458-463.
- Russell TA (2004). Comparison of sensory properties of whey and soy protein concentrates and isolates. A thesis of North Carolina State University, USA. MSc. Degree.
- Samanta SK (2013). Marine fish-derived bioactive peptides and proteins for human therapeutics. Int. J. Pharm. Pharmaceut. Sci. 5(3):132-146.
- San SC, Hyo KL, Chi WH, Eun SS, Chang YY, Myong JK, Na YK (2008). Physicochemical Properties of Isolated Peptides from Hwangtae (yellowish dlied pollack) Protein Hydrolysate. J. Food Sci. Nutri. 13:204-211.
- Sathe SK, Deshpande SS, Salunkhe DK (1982). Functional properties of winged bean (*Psophocarpus tetragonolobus, L*) proteins. J. Food Sci. 47: 503-506.
- Sikorski ZE (2002). Chemical and Functional Properties of Food Components. Second Edition. CRC Press LLC.
- Swaiswood HE (1996). Characteristics of milk, in: O.R. Fennema (Ed). Food Chemistry, Marcel Dekker, New York, pp. 841.
- Taheri A, Anvar SAA, Ahari H, Fogliano V (2012). Comparison the functional properties of protein Hydrolysates from poultry byproducts and rainbow trout (*Onchorhynchus mykiss*) viscera. Iranian Journal of Fisheries Sciences 12(1): 154-169.
- Wierenga PA, Gruppen H (2010). New views on foams from protein solutions. Current Opinion in Colloid and Interface Science 15(5):365-373.
- Wilde PJ, Clark DC (1996). The competitive displacement of blactoglobulin by Tween 20 from oil-water and air-water interfaces. J. Colloid and Interface Sci. 155:48-54.
- WR Egbert (2004). Isolated soy protein: Technology, properties, and applications, in: K.S. Liu (Ed.). Soybeans as Functional Foods and Ingredients, AOCS Press, Champaign, United States, pp. 134.